

Risk category, recommended monitoring and management for exposed HCW

Source
(close contact with positive COVID-19 patient)

Source with 3-ply surgical mask
(good control)

Source without 3-ply surgical mask
(no source control)

Risk to HCW

Risk to HCW

- No PPE
- Not wearing a 3-ply surgical mask

- Wearing a 3-ply surgical mask without eye protection/face shield
- Wearing a 3-ply surgical mask with eye protection/face shield but without gown and/or gloves

- No PPE
- Not wearing a 3-ply surgical mask
- Not wearing respirator (N95/PAPR) when performing AGP

- Wearing a 3-ply surgical mask without eye protection/face shield

- Wearing a 3-ply surgical mask with eye protection/face shield but without gown and/or gloves

MEDIUM

LOW

HIGH

MEDIUM

LOW

ACTIVE FOLLOW-UP

PASSIVE FOLLOW-UP

ACTIVE FOLLOW-UP

PASSIVE FOLLOW-UP

ASYMPTOMATIC

- Exclude from work for at least 14 days with home surveillance order and home assessment tool.
- Combined NP and OP swab for RT-PCR to be collected 72 hours after the potential exposure
- A repeat test after 48 hours must be carried out if the first test is negative
- If both initial tests are negative, a repeat NP and OP swab for RT-PCR test shall be performed at day 13 from date of exposure.

SYMPTOMATIC

- Exclude from work for at least 14 days with home surveillance order and home assessment tool.
- Combined NP and OP swab for RT-PCR to be collected immediately
- A repeat test after 48 hours must be carried out if the first test is negative
- If both initial tests are negative, a repeat NP and OP swab for RT-PCR test shall be performed at day 13 from symptom onset.

ASYMPTOMATIC

No testing or work restriction required

SYMPTOMATIC

No testing or work restriction required

ASYMPTOMATIC

- Combined Exclude from work for at least 14 days with home surveillance order and home assessment tool.
- Combined NP and OP swab for RT-PCR to be collected 72 hours after the potential exposure
- A repeat test after 48 hours must be carried out if the first test is negative
- If both initial tests are negative, a repeat NP and OP swab for RT-PCR test shall be performed at day 13 from date of exposure.

SYMPTOMATIC

- Exclude from work for at least 14 days with home surveillance order and home assessment tool.
- Combined NP and OP swab for RT-PCR to be collected immediately
- A repeat test after 48 hours must be carried out if the first test is negative
- If both initial tests are negative, a repeat NP and OP swab for RT-PCR test shall be performed at day 13 from symptom onset

ASYMPTOMATIC

No testing or work restriction required

SYMPTOMATIC

- Combined NP and OP swab for RT-PCR to be collected immediately
- If symptoms are persistent or worsening by 48-72 hours, test should be repeated immediately
- Exclude from work with MC and home assessment tool until symptoms resolve and negative 2nd test