UNIVERSITY MALAYA MEDICAL CENTRE GUIDELINE ON THE MANAGEMENT OF VACCINATED HEALTHCARE WORKERS

EXPOSED TO INDIVIDUALS WITH COVID-19 IN A HOSPITAL

(Prolonged Close Contact: Exposure at distance <1 metre and duration of >15 minutes)

COVID-19 PATIENT* YOU Vaccinated EXPOSURE RISK ACTION (SYMPTOMATIC)

ACTION (ASYMPTOMATIC)





HIGH RISK Quarantine for 10 days from date of exposure.

COVID-19 swab test immediately and on day 8 of exposure.

Quarantine for 10 days from date of exposure.

COVID-19 swab test immediately and on day 8 of exposure.

AGP PROCEDURE



NO MASK



NO MASK

SURGICAL MASK DID NOT USE N-95 OR FULL PPE

HIGH RISK

Quarantine for 10 days from date of exposure.

COVID-19 swab test immediately and on day 8 of exposure.

Quarantine for 10 days from date of exposure.

COVID-19 swab test immediately and on day 8 of exposure.

PATIENT COUGH INFRONT OF YOUR FACE



NO MASK



SURGICAL MASK DID NOT USE N-95 OR FULL PPE

HIGH RISK

Quarantine for 10 days from date of exposure.

COVID-19 swab test immediately and on day 8 of exposure.

Quarantine for 10 days from date of exposure.

COVID-19 swab test immediately and on day 8 of exposure.



NO MASK



SURGICAL MASK

MEDIUM RISK Do not need Quarantine.

COVID-19 swab test immediately and on day 8 of exposure.

Do not need Quarantine.

COVID-19 swab test immediately and on day 8 of exposure.

For healthcare workers who is under quarantine (7 or 10 days) must remain in quarantine for the duration of the quarantine although the initial COVID-19 swab test is negative (not-detected).

For healthcare workers with symptoms and was advice for immediate COVID-19 swab can go to the Special Staff Health Clinic during office hours or to Trauma and Emergency Department after office hours for assessment.

For healthcare workers who has a positive COVID-19 swab test, they will be contacted by the medical team for further management.

*COVID-19 patients include patients in wards and clinics, and other health workers who have a positive COVID-19 test.

UNIVERSITY MALAYA MEDICAL CENTRE GUIDELINE ON THE MANAGEMENT OF VACCINATED HEALTHCARE WORKERS

EXPOSED TO INDIVIDUALS WITH COVID-19 IN A HOSPITAL

(Prolonged Close Contact: Exposure at distance <1 metre and duration of >15 minutes)

COVID-19 PATIENT* YOU Vaccinated EXPOSURE RISK ACTION (SYMPTOMATIC)

ACTION (ASYMPTOMATIC)



MASK



NO MASK

MEDIUM RISK Do not need Quarantine.

COVID-19 swab test immediately and on day 8 of exposure.

Do not need Quarantine.

COVID-19 swab test immediately and on day 8 of exposure.



MASK



SURGICAL MASK



Do not need Quarantine.

COVID-19 swab test immediately.

Continue to work with strict adherence to the SOP.

COVID-19 swab test only if there are symptoms.



NO MASK



SURGICAL MASK AND FACE SHIELD



Do not need Quarantine.

COVID-19 swab test immediately.

Continue to work with strict adherence to the SOP.

COVID-19 swab test only if there are symptoms.

TAKING SWAB FOR COVID-19 OR AGP PROCEDURE



NO MASK



FULL PPE



Do not need Quarantine.

COVID-19 swab test immediately.

Continue to work with strict adherence to the SOP.

COVID-19 swab test only if there are symptoms.

For healthcare workers with symptoms and was advice for immediate COVID-19 swab can go to the Special Staff Health Clinic during office hours or to Trauma and Emergency Department after office hours for assessment.

For healthcare workers who have low risk and develop symptoms of COVID-19 will be given medical leave (MC) instead of home quarantine order. The duration of the MC will be for 2 days or until the result of the COVID-19 swab is available.

For healthcare workers who has a positive COVID-19 swab test, they will be contacted by the medical team for further management.

*COVID-19 patients include patients in wards and clinics, and other health workers who have a positive COVID-19 test.